



New findings from terrorism data: Dirichlet process random-effects models for latent groups

Minjung Kyung and Jeff Gill

Washington University, St Louis, USA

and George Casella

University of Florida, Gainesville, USA

[Received March 2010. Final revision March 2011]

Summary. Data obtained describing terrorist events are particularly difficult to analyse, owing to the many problems that are associated with the data collection process, the inherent variability in the data themselves and the usually poor level of measurement coming from observing political actors who seek *not* to provide reliable data on their activities. Thus, there is a need for sophisticated modelling to obtain reasonable inferences from these data. Here we develop a logistic random-effects specification using a Dirichlet process to model the random effects. We first look at how such a model can best be implemented, and then we use the model to analyse terrorism data. We see that the richer Dirichlet process random-effects model, compared with a normal random-effects model, can remove more of the underlying variability from the data, uncovering latent information that would not otherwise have been revealed.

Keywords: Empirical studies of terrorism; Generalized linear mixed models; Gibbs sampling; Hierarchical models; Logistic regression; Metropolis–Hastings algorithm; Terrorists

1. Introduction

The analysis of data on terrorists and terrorist attacks is problematic. These data are either observed public events, which omit planned but failed or cancelled attacks, or classified information at government agencies that are not available to general researchers. Also, most of the data sets focus purely on *incidents*: an observed violent attack along with covariates such as responsible group and target characteristics, as well as the extent of casualties and damage. This selection along with the outcome of interest, observable violent actions, effectively make the associated analytical work a *case–control study*.

Analysing terrorism incidents data has challenges that go significantly beyond those with regular events-based data or survey research data. Social science data can be extremely difficult to assess if the creators of the data set are unco-operative subjects. The key problem is that humans in covert, especially dedicated terrorist networks, for strategic reasons work to conceal not just their identities and intentions, but also their interactions with others. As a result of these collection issues, the data contain confounding effects, overlapping explanatory variables, high measurement error and unmeasured clustering forces. Confounding effects and overlapping explanations exist because the observed variables are often surrogates or summaries of the actual causal factors that are involved. The level of measurement here is typically from highly

Address for correspondence: George Casella, Department of Statistics, University of Florida, Gainesville, FL 32611, USA.
E-mail: casella@ufl.edu

non-granular nominal variables, and sometimes only purely qualitative information in the form of textual description. These problems could be considered missing data issues, but the general statistical tools for handling missing data (e.g. Little and Rubin (2002)) are unhelpful since the pattern of missingness here is controlled by actors who routinely seek to hide information.

Despite these seemingly insurmountable challenges, terrorism remains an extremely important problem because it affects personal safety, internal government policies, public perception and relations between nations. Normally, researchers who are faced with such problems would simply abandon the data at hand and develop strategies for collecting new, and more reliable, information. This simply is not feasible. Collecting data on terrorist activities beyond journalistic accounts is expensive, reliant on governments and sometimes even physically dangerous to the researcher. Although confounding, correlation and measurement error in explanatory variables can often be handled with careful model specification, the most daunting problem remains the latent clustering that occurs because terrorist actors are imitative of each other, communicate between groups and sometimes rely on the same resources.

We address the latent clustering problem with an improved implementation of non-parametric Bayesian generalized linear models using a logistic link function and Dirichlet process random effects. We have found that using Dirichlet process priors for the random effects can capture latent information in the data that would not otherwise be modelled (Gill and Casella, 2009). This provides a means of incorporating subclustering information in the data on terrorist events that would not otherwise be revealed in a standard context, even if it cannot be directly parameterized in the model, thus improving model fit. The term ‘subcluster’ is used in this context instead of ‘cluster’ since the modelling approach does not penalize for overfragmentation of the real clusters that are hidden in the data. Here we use this tool to study suicide terrorist attacks in the Middle East and northern Africa from 1998 to 2004.

Our approach is based on a new Gibbs sampling method that was developed in Kyung *et al.* (2010a). In the work here, we generalize the newly developed Gibbs sampling based on a mixture representation of the logistic distribution in Kyung *et al.* (2010a), and we find that the Gibbs sampler appears to mix more rapidly than the commonly used alternative slice sampler of Damien *et al.* (1999). We show that the slice sampler typically has higher auto-correlation and poorer mixing than the approach that is taken here.

Our results support a theorized but not fully explored notion about terrorist groups. There are groups or clusters of terrorists who work in similar fashion, either because they communicate or because they emulate each other. Information from the Markov chain Monte Carlo sampling scheme produces coefficient estimates that account for this latent trait and therefore condition on it. Our findings demonstrate that there are reliable coefficient estimates for attack characteristics that are shared across groups in the Middle East and northern Africa over the period 1998–2004. This includes some sensationalist attacks against large government buildings, as well many more routine smaller events.

2. Background on terrorism data sets

There are now several major databases on terrorist incidents stored at academic institutions like the University of Maryland (START) and at government agencies like the US Homeland Security Agency. Event data on terrorist hostage incidents are provided by ‘International terrorism: attributes of terrorist events’, which was originally assembled by Mickolus (1982) and later updated (Mickolus, 2006). This database records transnational terrorist incidents and therefore ignores so-called ‘domestic terrorism’. Conversely the data set ‘Political violence in the United States, 1819–1968’ (Levy *et al.*, 1969) lists incidents and covariates for political violence resulting

in injury or death for about 150 years. The International Policy Institute for Counter-Terrorism in Herzlia, Israel, provides a data set of terrorist attacks in Israel. The US Department of Homeland Security supports the National Memorial Institute for the Prevention of Terrorism terrorism knowledge base, which provides on line a listing of terrorism incidents with information on the terrorists and an emphasis on legal information. Another on-line listing is the global terrorism database which includes information on global terrorist events starting from 1970. Researchers have used these primarily for creating summary statistics and basic tabular analyses. Standard statistical modelling has yielded some insights into the determinants and timing of terrorists incidents (Enders (2007), Enders and Sandler (1995) and Li and Schaub (2004), for example), but with limited results.

Game theoretic approaches have been used to circumvent the problem of data quality. The way that governments and terrorists (factions) strategize has been studied, for instance, to explain why extremist groups often increase terrorist activity after a government has made concessions to moderate factions. These results have yielded important insights about terrorism in Ireland, Spain and other countries (Bueno de Mesquita (2005); see also such works as Bueno de Mesquita and Dickson (2007), Arce and Sandler (2007), Sandler and Arce (2003), Kydd and Walter (2002) and Siqueira and Sandler (2006)), but they are not in general motivated by conventional data analysis.

Another approach is to build models of networks of terrorist and terrorist organizations. Despite the convention that terrorists should not be treated as unitary actors (Chai, 1993; Crenshaw, 1981), the study of terrorist organizations as networks is less developed. Social network analysts have discovered that covert organizations tend to be cellular and distributed rather than hierarchical (Carley, 2004; Krebs, 2002; Rothenberg, 2002), and the government has supported research to model these as standard networks through the National Research Council. Carley (2003, 2006) used 'meta-matrices' that capture not just the ties between terrorists but also their knowledge and tasks from semi-automatic parsing of signal traffic (Tsvetovat and Carley, 2006). This work yields, among other things, lessons about how best to 'destabilize' terrorist networks (Tsvetovat and Carley, 2005; Moon and Carley, 2007).

None of these research tracks has been very successful in building standard semicausal regression models that are preferred in the social sciences. This is because the data are, in general, poorly measured and highly non-granular. Specifically, we observe roughly measured categorical variables produced by governments as they record and react to attacks on their soil or against their citizens elsewhere. A huge part of the problem is that the primary actors under study are deliberately trying to prevent such data from being collected in an accurate and useful manner. So, unlike many missing data or poorly measured data problems, the key agents in the data generation process are not denying observation in benign ways; they are attempting to deny observation in wilful and strategic ways. Key information, that is almost always missing from such data, is the intentions of terrorists, strategic alternatives that they face and actions that failed to achieve any result. All of these problems above mean that the analyst has a doubly difficult task in creating meaningful inferential models.

3. Data on suicide attacks

The data that we use here come from the global terrorism database II (LaFree and Dugan, 2008), restricted to events in the Middle East and northern Africa from 1998 to 2004. After removing almost totally incomplete cases, this provides 1041 violent attacks by terrorist groups, 154 (15%) of which were suicide attacks where at least one of the individual assailants was killed by design. Our outcome variable of interest is therefore the dichotomous observation of a suicide attack

or not. Suicide attacks pose a substantially higher challenge for governments since the assailant has great control over placement and timing and also does not need to plan his or her escape (Pape, 2006). Thus they may inflict severe damage to otherwise safe targets.

Terrorism data are notoriously difficult to model from a regression context, so we first pick a rich set of explanatory variables. Time is important during this era. It is also complex. Our approach is to designate the start of the period of study as an indicator variable so that estimated positive or negative coefficients are revealing about a trend. There were 273 terrorist attacks worldwide in 1998 with a recorded high of 741 killed along with 5952 injured (US State Department: <http://www.state.gov/www/global/terrorism/1998report/1998index.htm>). Of these 273, 103 were in our geographic area of study. This particular year was also notable for the incredibly destructive simultaneous bombings of the US Embassies in Nairobi, Kenya (291 killed; roughly 5000 injured), and Dar es Salaam, Tanzania (10 killed; 77 injured), in August.

Relatedly, we also consider a variable, *MULT.INCIDENT*, which indicates whether the attack is part of a co-ordinated multisite event. This was so in a surprisingly large number of cases, 136 (13.1%). Including this explanatory variable in the model is a means of testing the relationship between connected events at different sites and the use of suicide attackers.

Sometimes multiple parties are involved in a regional conflict, such as the wars in Iraq and Afghanistan. This can make it more difficult to assess responsibilities. Strategic actors sometimes make false claims about responsibility (both for and against) as a means of concealing actions and intentions. Normally, but not always, a suicide attack can be traced to a specific group since identifying biological information is left at the incident site. Also, in some settings, the family is forthcoming about the identity of the attacker. We measure this effect with the variable *MULT.PARTY*, where (also) 136 out of 1041 cases are coded as 1.

There can also be terrorist incidents in which there is substantial uncertainty about the identity of any single attacking group. The variable *SUSP.UNCONFIRM* is coded as 1 (209/1041) if government officials express notable doubt about attributing responsibility. As with the involvement with multiple parties, we expect this to be less common for suicide attacks (25/209, which is 12% of such cases).

One key issue is the linkage between the success of the attack and the suicide nature of the attack. Obviously, there is some linkage or groups would not persist in organizing suicide attacks over time. Since this is not in question, we ask the question from the perspective of the attacked party, whose law enforcement, military and intelligence services seek to understand and prevent further attacks. By using the variable *SUCCESSFUL* (which was true in 966 of the events) on the right-hand side of the model specification, we are asking: given that it is a successful attack, how likely is it that a suicide assailant was used?

An important issue is the type of attack that is used. The data are presented in a quasi-ordered fashion with the categories (with counts in parentheses) armed assault (385), assassination (47), bombing or explosion (514), facility or infrastructure attack (50), hijacking (7), barricade incident (3) and kidnapping (35). The model results were highly robust to different recodings of this variable since armed assaults and bombings dominate. We therefore left the original *ATTACK.TYPE* coding in place such that for suicide attackers a positive estimated coefficient shows a preference for bombing strategies and a negative estimated coefficient shows a preference for armed assault. We also consider a variable, *WEAPON.TYPE*, that is coded 1 for the use of explosives, dynamite, or general bombs to help to differentiate the suicide bombings from non-suicide bombings (Table 1). This relationship is unbalanced since the bulk of attacks are not from suicide attackers. However, suicide attackers are typically more effective in creating casualties and the psychology of suicide attackers is more effective in creating terror

Table 1. Suicide *versus* general bomb attacks

	<i>Suicide versus bomb</i>	
	<i>Not bomb</i>	<i>Bomb</i>
Not suicide	720	661
Suicide	5	224

within populations. The key relationship here is the heavy use of bombs by suicide attackers. Note that our subsequent analysis uses the coding that is described in `ATTACK.TYPE`, not the dichotomization in Table 1.

Of course the targets that terrorists pick are important to understand. The primary distinction, which we focus on here, is civilian (832 cases; 0 coding) *versus* military (209 cases; 1 coding) targets. As noted terrorists tend to prefer civilian targets since they are usually ‘softer’ (less guarded) and more plentiful, and the perceived randomness of death and destruction has a greater effect on the morale of the citizens. This explanatory variable is labelled `TARGET.TYPE`.

In addition to being successful, we also want to incorporate the extent of human damage that the terrorist attack inflicts. Again, we look at this not from the terrorist perspective but from the government perspective by including `NUM.FATAL` (3424 total) and `NUM.INJUR` (8123 total) on the right-hand side of the model. So our substantive question is, for higher (or lower) levels of casualties, what is the likelihood, controlling for other factors, that the terrorists used a suicide attacker? Another measure of damage is the negative psychological or social effect that terrorist attacks inflict on national populations. In fact, this is usually the most widespread effect that such events produce. The collectors of these data subjectively coded this into `PSYCHOSOCIAL` with ascending levels: none, minor, moderate and major (a summary is given below). We also look at a variable for property damage, which can vary substantially. The data are provided with yes–no values along with ‘minor’ in between in the variable `PROPERTY.DAMAGE`.

3.1. Visualizing the suicide data

Consider the four coplots in the four frames of Fig. 1, where two variables are plotted in each, at levels of a different binary third variable, with different mixes of variables.

Note the similarities and differences in the distribution of fatalities between Figs 1(a) and 1(b). The annual patterns of fatalities for not using a bomb and attacking civilian targets are very close, except for some high death events at the end of the series in Fig. 1(a), which are attributable to bombs used on non-military targets. Thus the relationship is subtle and it spans four variables: year, military target, bomb used and number of fatalities. This confounding of effects leads to collinearity and is typical of terrorism data. Substantively, the key is that military targets are more hardened towards terrorist attacks, even if they are possibly more desirable political targets by some groups. Civilian attacks also dominate in the data, 867 to 174. Since terrorism data are difficult to obtain, it is tempting to use as many variables as one can reasonably obtain on the right-hand side of regression models. Yet the observed collinearity problem makes this problematic. So we would like to use `TARGET.TYPE` and `WEAPON.TYPE` as explanatory variables since they reflect a graphically observable difference in effect. However, this is deleterious to model in conventional ways, which is why we develop a Dirichlet process for random-effects models which accounts for unobserved clumping in the data that comes from these overlapping effects.




Fig. 1. Coplots for Middle East and northern Africa suicide attacks, 1998–2004: (a), (b) comparison of the year starting with 1998 in the study (x-axis) with the number of fatalities for these years (y-axis) (the comparison is split by whether (a) the target was civilian or military (control, TARGE.TYPE) and whether (b) a bomb was used (control, WEAPON.TYPE)); (c), (d) comparison of the levels of psychosocial effect (x-axis) against the number injured (the comparison is split whether (c) there were multiple associated incidents for the single attack (control, MULTI.INCIDENT) and by whether or not (d) the government attacked has suspicions about the attackers' identity (control, SUSP.UNCONFIRM))

Table 2. Summary for PSYCHOSOCIAL

Level	none	minor	moderate	major
Count	18	946	66	11

The real underlying goal for terrorists is not actually the results of the physical violence that is enacted. Rather it is undermining citizens' confidence in their government's ability to protect them. Thus gory attacks against easy targets that receive widespread media coverage are considerable victories for the planners of such attacks. To explore this strategic consideration we focus on the relationship between escalating negative psychological or social impact and the number injured. Injured people are a fundamentally different kind of casualty effect from fatalities because they wander around the scene of the attack, command medical attention, agree to media interviews and spend the rest of their life discussing the event with friends and relatives. Thus although fatalities provide horrific statistics, injuries take on a longer psychological life among citizens of the attacked nation. Therefore Figs 1(c) and 1(d) plot the intensity of the psychological effect (x -axis) against the magnitude of the number of injuries (y -axis).

In Fig. 1(c) it is easy to see a distinctly different pattern between PSYCHOSOCIAL and NUM.INJUR for levels of the third dichotomizing variable: whether or not the attack contained multiple associated events (MULT.INCIDENT). Table 2 shows that the bulk of the psychological or social effects are minor, and these are concentrated in the 'No' part of Fig. 1(c) with low levels of injuries. In contrast, the measures of psychological or social impact for multiple incidents are almost all minor and moderate but show higher average injuries. The third dichotomizing variable in Fig. 1(d) is whether or not the government has notable doubts about the identity of the attacking party. Here we see that injuries are generally higher when there is certainty about the attacking group, but also with greater variance. Difficult-to-identify groups may inflict less damage because they are less bold in their planning as a means of reducing the government's detective ability. Figs 1(c) and 1(d) show the same challenge as Figs 1(a) and 1(b): there are interesting, and perhaps important, features in the control variables, but the high level of collinearity makes these difficult or impossible to include in the same model as covariates.

This exploratory graphical analysis shows that data on terrorist attacks provide some special challenges. First, the level of measurement is usually low, with mostly categorical, and some purely qualitative, variables. Second, as shown in Fig. 1, there are collinear relationships in combinations of variables at specific values that can easily be averaged over, and therefore missed, by standard regression style models. Third, correlation between these variables is almost certainly due to additional factors. Finally, these data sets are almost always missing key variables about the *intentions*, *strategies* and even *failings* of the terrorist groups. If these groups are heterogeneous in such respects, and in their effectiveness, then we would expect subtle distinctions as seen in Fig. 1. So it is this appreciation for the difficulties that are inherent in the data analytic understanding of terrorism that moves us to describing a Bayesian non-parametric set-up that is intended to make finer grouping distinctions through Dirichlet priors on random effects to capture latent variability.

4. Generalized linear models and random effects

We need a model structure that accounts for the problems with data that were discussed in

Section 3, particularly the issue of latent groupings based on unobserved relationships between cases. Although it is impossible to provide a reliable coefficient estimate of the magnitude of these effects in the general regression modelling sense since there are no overt indicators, we can develop a model that *accounts for* this phenomenon and thus fits such data better than alternatives that ignore latent groupings. In this section we start with conventional non-linear regression modelling and build up to our full non-parametric Bayes specification for this purpose.

4.1. Generalized linear models

Generalized linear models have enjoyed considerable attention over the years, providing a flexible framework for modelling discrete responses by using a variety of error structures. If we have observations that are discrete or categorical, $\mathbf{y} = (\mathbf{y}_1, \dots, \mathbf{y}_n)$, such data can often be assumed to be independent and from a distribution in the exponential family, where the likelihood function has components of the model such as the form of the link function and the type of error structures that result. The classic book by McCullagh and Nelder (1989) describes these models in detail; see also the developments in Dey *et al.* (2000) or Fahrmeir and Tutz (2001).

4.2. Generalized linear mixed models

A generalized linear *mixed* model is an extension of a generalized linear model that allows random effects and can give us flexibility in developing a more suitable model when the observations are correlated, or where there may be other underlying phenomena that contribute to the resulting variability. Thus, the generalized linear mixed model can be specified to accommodate outcome variables conditional on mixtures of possibly correlated random and fixed effects (Breslow and Clayton, 1993; Buonaccorsi, 1996; Wang *et al.*, 1998; Wolfinger and O'Connell, 1993). Details of such models, covering both statistical inferences and computational methods, can be found in McCulloch and Searle (2001) and Jiang (2007).

4.3. Dirichlet process random effects

Dirichlet process mixture models were introduced by Ferguson (1973) and Antoniak (1974) and have been the subject of much research since then, most of which described the theoretical character of the process, and some estimation strategies (notably Blackwell and McQueen (1973)). In current applications, a popular strategy is to use Dirichlet process priors in hierarchical models. Work by Escobar and West (1995), MacEachern and Müller (1998), Neal (2000) and Teh *et al.* (2006) developed models and Markov chain Monte Carlo schemes to fit such models.

Recently, Dorazio *et al.* (2007) used a generalized linear mixed Dirichlet model (GLMDM) with a log-link for spatial heterogeneity in animal abundance. They proposed an empirical Bayesian approach with the Dirichlet process, instead of the regular assumption of normally distributed random effects, because they argued that, for some species, the sources of heterogeneity in abundance are poorly understood or unobservable. They noted that the Dirichlet process prior is robust to errors in model specification and allows spatial heterogeneity in abundance to be specified in a data-adaptive way. Gill and Casella (2009) suggested a GLMDM with an ordered probit link to model political science data, specifically modelling the stress, from public service, of Senate-confirmed political appointees in the USA as a reason for their short tenure. For the analysis, a semiparametric Bayesian approach was adopted, using the Dirichlet process for the random effect.

More recently Kyung *et al.* (2010b) developed algorithms for estimation of the precision parameter and new Markov chain Monte Carlo algorithms for a linear mixed Dirichlet random-effects models. Also, they showed how to extend the results to a generalized Dirichlet process mixed model with a probit link function and used a new parameterization of the hierarchical model to derive a Gibbs sampler that more fully exploits the structure of the model and mixes very well. They could also establish that the sampler proposed is an improvement, in terms of operator norm and efficiency, over other commonly used algorithms. Kyung *et al.* (2010a) extended the available sampling schemes to handle other link functions, including logistic and log-linear. Here we apply their logistic sampling scheme to fit our terrorism data.

5. Logistic mixed Dirichlet process models

Let \mathbf{X}_i be covariates associated with the i th observation, β be the coefficient vector and ψ_i be a random effect accounting for subject-specific deviation from the underlying model. Assume that the $Y_i|\psi$ are conditionally independent, each with a density from the exponential family, where $\psi = (\psi_1, \dots, \psi_n)$. Then, based on the generalized linear mixed model notation in McCulloch and Searle (2001), the GLMDM can be expressed as follows. Start with the generalized linear model that was described above,

$$\begin{aligned}
 Y_i|\gamma &\stackrel{\text{ind}}{\sim} f_{Y_i|\gamma}(y_i|\gamma), & i = 1, \dots, n, \\
 f_{Y_i|\gamma}(y_i|\gamma) &= \exp\{\{y_i\gamma_i - b(\gamma_i)\}/\xi^2 - c(y_i, \xi)\}, & (1)
 \end{aligned}$$

where y_i is discrete valued. Here, we know that $E[Y_i|\gamma] = \mu_i = \partial b(\gamma_i)/\partial \gamma_i$. Using a link function $g(\cdot)$, we can express the transformed mean of Y_i , $E[Y_i|\gamma]$, as a linear function, and we add a random effect to create the mixed model:

$$g(\mu_i) = \mathbf{X}_i\beta + \psi_i. \tag{2}$$

Here, for the Dirichlet process mixture models, we assume further that

$$\psi_i \sim G, \quad G \sim \mathcal{DP}(mG_0), \tag{3}$$

where \mathcal{DP} is the Dirichlet process with base measure G_0 and precision parameter m . The base measure functions as an expected distribution in this set-up, and the precision parameter determines the variability around this base measure, which is sometimes called the ‘cylinder’ around it as a visual image. The base measure can even be specified as a normal distribution, meaning that m determines the variability of the distributions around this choice. With this model, we relax the normal assumption, and we provide a richer model to capture more variabilities in the random effects.

Blackwell and MacQueen (1973) proved that, for ψ_1, \dots, ψ_n independent and identically distributed (IID) from $G \sim \mathcal{DP}$, the joint distribution of ψ is a product of successive conditional distributions of the form

$$\psi_i|\psi_1, \dots, \psi_{i-1}, m \sim \frac{m}{i-1+m}g_0(\psi_i) + \frac{1}{i-1+m} \sum_{l=1}^{i-1} \delta(\psi_l = \psi_i) \tag{4}$$

where $\delta(\cdot)$ denotes the Dirac δ -function and $g_0(\cdot)$ is the density function of base measure. Thus, with n observations from a Dirichlet process with precision parameter m , the marginal distribution of a partition $\{n_1, n_2, \dots, n_k\}$, where $\sum_j n_j = n, n_j \geq 1$, is given by

$$\pi(n_1, n_2, \dots, n_k) = \frac{\Gamma(m)}{\Gamma(m+n)} m^k \prod_{j=1}^k \Gamma(n_j), \tag{5}$$

which is a normalized probability distribution on the set of all partitions of n observations.

We now define a *partition* C to be a grouping of the sample of size n into k groups, $k = 1, \dots, n$, and we call these subclusters since the grouping is done non-parametrically rather than on substantive criteria, i.e. the partition assigns different parameters across groups and the same parameters within groups; cases are IID only if they are assigned to the same subcluster. Furthermore, the use of ‘subcluster’ is important to distinguish this grouping from substantive clusters since there is no penalty term in the model fit for increasing their number.

Applying Lo (1984), lemma 2, and Liu (1996), theorem 1, to formula (4), we can calculate the likelihood function, which by definition is integrated over the random effects, as

$$L(\theta|\mathbf{y}) = \frac{\Gamma(m)}{\Gamma(m+n)} \sum_{k=1}^n m^k \sum_{C:|C|=k} \prod_{j=1}^k \Gamma(n_j) \int f(\mathbf{y}_{(j)}|\theta, \psi_j) dG_0(\psi_j),$$

where C defines the subclusters, $\mathbf{y}_{(j)}$ is the vector of y_i s that are in subcluster j and ψ_j is the common parameter for that subcluster. There are $S_{n,k}$ different subclusters C , the Stirling number of the second kind (Abramowitz and Stegun (1972), pages 824–825).

The partition C can be represented by an $n \times k$ matrix \mathbf{A} defined by

$$\mathbf{A} = (a_1, a_2, \dots, a_n)'$$

where each a_i is a $1 \times k$ vector of all 0s except for a 1 in the position indicating which group the observation is from. Thus, \mathbf{A} represents a partition of the sample of size n into k groups, with the column sums giving the subcluster sizes. Note that both the dimension k and the placement of the 1s are random, representing the subclustering process. On the basis of the representation in McCullagh and Yang (2006), if the partition C has subclusters $\{S_1, \dots, S_k\}$, then, if $i \in S_j$, $\psi_i = \eta_j$ and the random effect can be rewritten as

$$\psi = \mathbf{A}\boldsymbol{\eta}, \tag{6}$$

where $\boldsymbol{\eta} = (\eta_1, \dots, \eta_k)$ and $\eta_j \stackrel{\text{IID}}{\sim} G_0$ for $j = 1, \dots, k$.

Usually, in a regular generalized linear mixed model the random effects accounting for subject-specific deviation from the underlying model, ψ s, are assumed to be distributed as $N(0, \sigma_\psi^2)$. In this paper, we assume that $\psi_i \sim \mathcal{DP}\{m, N(0, \tau^2)\}$, $i = 1, \dots, n$, independent. Thus, for $\boldsymbol{\eta}$, we obtain $\boldsymbol{\eta} \sim N_k(0, \tau^2 I)$. These ψ s are random effects to capture the issue of latent groupings. Because of the properties of the Dirichlet process that we discussed above, we can develop a rich model with the random effects for the unobserved relationships.

In this particular paper, we consider only models for the binary responses with a logit link function. So we model

$$Y_i \sim \text{Bernoulli}(p_i), \quad i = 1, \dots, n,$$

where y_i is 1 or 0, and $p_i = E[Y_i]$ is the probability of a success for the i th observation, and the sampling distribution is

$$f(\mathbf{y}|\mathbf{A}) = \int \prod_{i=1}^n \left[\frac{\exp\{\mathbf{X}_i\boldsymbol{\beta} + (\mathbf{A}\boldsymbol{\eta})_i\}}{1 + \exp\{\mathbf{X}_i\boldsymbol{\beta} + (\mathbf{A}\boldsymbol{\eta})_i\}} \right]^{y_i} \left[\frac{1}{1 + \exp\{\mathbf{X}_i\boldsymbol{\beta} + (\mathbf{A}\boldsymbol{\eta})_i\}} \right]^{1-y_i} dG_0(\boldsymbol{\eta}), \tag{7}$$

which typically can only be evaluated numerically. Here the general link function is

$$p_i = g^{-1}\{\mathbf{X}_i\boldsymbol{\beta} + (\mathbf{A}\boldsymbol{\eta})_i\} = [1 + \exp\{-\mathbf{X}_i\boldsymbol{\beta} - (\mathbf{A}\boldsymbol{\eta})_i\}]^{-1}.$$

This model can now be fitted by using a variation of the Markov chain Monte Carlo scheme of Kyung *et al.* (2010b), where we use a mixture representation of the logistic distribution (Andrews and Mallow, 1974). Details are in Appendix A.

6. Simulation study

We evaluate our sampler through a simulation study. We need to generate outcomes from Bernoulli distributions with random effects that follow the Dirichlet process. To do this we fix K , the true number of subclusters (which is unknown in actual circumstances); then we set the parameter m according to the relation

$$K = \sum_{i=1}^n \frac{m}{m + i - 1}, \tag{8}$$

where we note that, even if \hat{m} is quite variable, there is less variability in $\hat{K} = \sum_{i=1}^n \hat{m} / (\hat{m} + i - 1)$. When we integrate over the Dirichlet process, as done algorithmically according to Blackwell and McQueen (1973), the right-hand side of equation (8) is the expected number of clusters, given the prior distribution on m . Neal (2000), page 252, showed this as the probability in the limit of a unique table seating, conditional on the previous table seatings, which makes intuitive sense since this expectation depends on individuals sitting at unique tables to start a new (sub)cluster in the algorithm.

Using the GLMDM with the logistic link function, we set the parameters: $n = 100$, $K = 40$, $\tau^2 = 1$ and $\beta = (1, 2, 3)$. Our Dirichlet process for the random effect has precision parameter m and base distribution $G_0 = N(0, \tau^2)$. Setting $K = 40$ yields $m = 24.21$. Independently generated X_1 and X_2 from $N(0, 1)$ are used as the fixed design matrix to generate the binary outcome Y . Then the Gibbs sampler was iterated 200 times to obtain values of m , A , β , τ^2 and η . This procedure was repeated 1000 times, saving the last 500 draws as simulations from the posterior.

We compare the slice sampler with the Gibbs sampler with the Kolmogorov–Smirnov (KS) distribution normal scale mixture with the prior distribution of β from $\beta | \sigma^2 \sim N(\mu \mathbf{1}, d^* \sigma^2 I)$ and a flat prior on μ , $\mu \sim \pi(\mu) \propto c$. For the estimation of K , we use the posterior mean of m , \hat{m} and calculate \hat{K} by using equation (8). The starting points of β come from the maximum likelihood estimates by using iteratively reweighted least squares. All summaries in the tables are posterior means and standard deviations calculated from the empirical draws of the chain in its apparent converged (stationary) distribution.

The numerical summary of this process is given in Table 3. The estimate of β with KS mixture sampler is closer to the true value than those with the slice sampler, with smaller standard deviation. To evaluate the convergence of β , we consider the auto-correlation function plots that are given in Fig. 2. The Gibbs sampler of β from the slice sampler exhibits strong auto-correlation, implying poor mixing. The estimates of K were 43.0423 with standarderror 4.2670 from the

Table 3. Estimation of the coefficients of the GLMDM with logistic link function and the estimate of K , with true values $K = 40$ and $\beta = (1, 2, 3)$ †

Estimation method	β_0	β_1	β_2	K
Slice sampler	2.2796 (0.4628)	3.2709 (0.5558)	4.7529 (0.7208)	43.0423 (4.2670)
KS mixture	0.4900 (0.2024)	1.0494 (0.2468)	1.7787 (0.2491)	43.4646 (4.0844)

†Standard errors are in parentheses.




Fig. 2. Auto-correlation function plots for the GLMDM with logistic link: (a) β_0 , slice sampler; (b) β_1 , slice sampler; (c) β_2 , slice sampler; (d) β_0 , KS mixture sampler; (e) β_1 , KS mixture sampler; (f) β_2 , KS mixture sampler

slice sampler and 43.4646 with standard error 4.0844 from the KS mixture sampler. Obviously these turned out to be good estimates of the true $K = 40$.

7. Application of the model to data on suicide attacks

We apply a GLMDM specification with a logit link function, as described, and for the model parameters with the prior distributions in expression (13) in Appendix A.2 where $\mathbf{B} = (\mu, 0, \dots, 0)'$, $d^* = 5$ and (a, b) are fixed as (3,2). We ran the Markov chain for 50000 iterations, disposing of the first half. There is no evidence of non-convergence in these runs by using standard diagnostic tools (e.g. with the commonly accepted procedures: Geweke, Gelman and Rubin, Heidelberger and Welch, and time series graphics; see Gill (2007), chapter 12). Table 4 provides results from two approaches: a standard Bayesian random-effects logit model (BRELM) with flat priors, and a Dirichlet random-effects model, with posterior means, posterior standard errors and 95% highest posterior density (HPD) intervals.

Note from Table 4 that, although there are no changes in sign or statistical reliability for the estimated *coefficients* (posterior means) between the two models, the magnitudes of the effects are uniformly smaller with the enhanced model and four of the BRELM coefficient estimates are almost twice as large as the corresponding GLMDM coefficient estimates. This indicates that there is extra information in the data that is detected by the Dirichlet random effect that tends to dampen the size of the effect of these explanatory variables on explaining incidences of terrorist attacks. Specifically, running the standard random-effects model would find an *exaggerated* relationship between these explanatory variables and the outcome. In other work (Gill and Casella, 2009), we observed that the GLMDM model provided a theoretically important subset of coefficient estimates that were greater in magnitude, so the non-parametric information does not always diminish the resulting effect sizes.

These results are interesting substantively. The year 1998 does stand out relatively speaking (until 2001 in the USA of course, but that case is not in our geographic focus). This is a confirmatory finding since we know that 1998 was exceptional from empirical and journalistic

Table 4. Global suicide attack data—coefficients, standard errors and HPD intervals from a BRELM and from the GLMDM using a logit link

Variable	Results from BRELM				Results from GLMDM–logit			
	Coefficient	Standard error	95% HPD		Coefficient	Standard error	95% HPD	
Intercept	−6.457	4.232	−21.605	−3.407	−4.105	0.559	−5.276	−3.079
YEAR—1998	0.303	0.228	0.135	1.137	0.195	0.039	0.121	0.273
MULT.INCIDENT	−0.802	0.488	−2.222	−0.142	−0.585	0.221	−1.028	−0.162
MULTI.PARTY	−0.945	0.690	−3.289	−0.225	−0.626	0.229	−1.088	−0.189
SUSP.UNCONFIRM	−0.109	0.344	−0.928	0.472	−0.061	0.198	−0.455	0.331
SUCCESSFUL	−1.035	0.705	−3.308	−0.262	−0.695	0.245	−1.172	−0.210
ATTACK.TYPE	0.122	0.135	−0.122	0.466	0.098	0.073	−0.046	0.240
WEAPON.TYPE	2.714	1.673	1.346	7.769	1.725	0.320	1.162	2.422
TARGET.TYPE	−0.073	0.330	−0.749	0.527	−0.038	0.185	−0.434	0.323
NUM.FATAL	−0.019	0.025	−0.085	0.017	−0.013	0.012	−0.036	0.009
NUM.INJUR	0.030	0.030	0.010	0.126	0.017	0.004	0.008	0.025
PSYCHOSOCIAL	0.824	0.633	0.216	3.044	0.555	0.192	0.188	0.944
PROPERTY.DAMAGE	0.439	0.305	0.122	1.406	0.297	0.094	0.114	0.483

sources. It also appears that multiple co-ordinated incidents are less associated with suicide attacks (again September 11th, 2001, being a notable exception) since MULT.INCIDENT is negative and statistically reliable. This is interesting and it suggests that planners of simultaneous linked terrorist events find it more difficult to manage multiple suicidal agents. MULTI.PARTY in Table 4 tells a similar story; co-ordinated groups of attackers are less likely to plan suicide attacks. This may be because together they have more traditional military resources at hand.

Interestingly, if the attack is successful it is less likely to be a suicide attack. When the delivery vehicle is a person, there are more variables to worry about, in particular whether the individual is sufficiently practised and indoctrinated. We know from qualitative accounts that fervent nationalism and religious extremism are critical components of this process. Unfortunately for the planners of such attacks, the efficacy of these means is almost certainly negatively correlated with the subject's age and intelligence.

With regard to type of attack and type of weapon, there is strong evidence that those plotting bomb attacks (generally speaking) are more likely to consider suicide agents to deliver the weapon. Note that the coefficient for WEAPON.TYPE is more statistically reliable than the coefficient for ATTACK.TYPE, although they agree on direction.

Increased damage to property is positively associated with a suicide rather than non-suicide attack. This appears to be related to the terrorists' preference for easier civilian targets where an equivalent-sized device will produce more damage than against better protected government targets.

In addition to ATTACK.TYPE, three other estimated coefficients do not appear to be reliable at standard thresholds in the resulting model: TARGET.TYPE, SUSP.UNCONFIRM and NUM.FATAL (their 95% HPD intervals cover zero). Since 20% of the types of target were military *versus* civilian, it is somewhat surprising not to find a reliable effect conditioning on the rest of the model. More interestingly, greater fatalities at the event site do not suggest a greater probability of the damage being produced by a suicide attacker. The apparent, but unproven, lack of a relationship here suggests that suicide attacks are more driven by other intended consequences like widespread injuries and the psychological or social effect, both of which show up here as coefficient estimates that have substantial statistical support. This finding, although surprising to some, is consistent with a large proportion of the literature on the intentions of terrorists. Terrorism is most fundamentally aimed at diminishing a national population's confidence in the ability of the government to defend them at home.

Finally, we also note that the GLMDM–logit model has uniformly smaller HPD intervals than the standard generalized linear mixed model–logit model, BRELM. This is consistent with the findings in Kyung *et al.* (2009) for models with other link functions. The comparison is shown dramatically in Fig. 3, where we see that the HPD intervals for the standard random-effects logit model (grey) are substantially wider than those for the new method (black). Thus, the richer random-effects model can remove more extraneous variability, providing tighter HPD intervals. This is evidence that the Dirichlet procedure is capturing additional latent information. So the non-parametric Bayes approach that is taken here provides more *conservative* estimates of effect sizes (smaller absolute coefficient estimates) *and* gives more accuracy for these estimates (smaller HPD intervals). Thus we have substantially improved the state of data analysis in an area where researchers struggle.

8. Discussion

In this paper we employ a new methodology to solve an old problem. Terrorism has existed since humans first built weapons, and the academic study of terrorism increased dramatically




Fig. 3. 95% HPD intervals: standard logit (grey bar, vertical line) versus GLMDM-logit (black bar, vertical line) (the break in the grey intercept bar indicates that the lower end is beyond the scale of the graph; vertical bars denote the posterior means from the two models for each coefficient)

after September 11th, 2001. In fact, a non-field-specific www.jstor.org search with the word 'terrorism' provides 26426 article citations ranging from the year 1848 to the year 2010. The first citation for the year 2002 occurs at number 16804, meaning that 36% of the 162-year history occurs after the 154th year (the last 5%). Although this is not a complete analysis, it clearly indicates a strong upswing after 2001.

Unfortunately, the data routinely present formidable challenges to analysis with conventional statistical tools, since such data are always more messy and interrelated than in other applications. The key underlying issue is that a set of diversely organized covert operatives do not co-operate with data collection efforts and the resulting product is often poorly measured.

Our methodological approach uses a variant of non-parametric Bayesian generalized linear models with a logistic link function and Dirichlet process random effects. We developed and verified the properties of new Markov chain Monte Carlo estimation tools for this problem in previous work, noting that they incorporate latent information such that models fit better in the presence of unexplained heterogeneity in the data. This turns out to be ideal for the empirical analysis of terrorism, which has diverse and subversive operators generating the heterogeneous data. There is evidence from Table 4 that the GLMDM–logit model's superior properties over the analogous standard BRELM come from fitting on the latent scale since that is the only fundamental distinction between the two. The inherent binning to subclusters, that is part of our sampler, reaches this improvement by recognizing and modelling this underlying heterogeneity.

What did we learn about Middle East and northern African terrorism? In general multiple groups working together do not focus on using suicide attackers. Instead, they appear to work in more quasi-military fashion with standard weapons. Surprisingly, the more damaging attacks were more likely to be from non-suicide assailants. This runs counter to media reports, which tend to focus on the goriness of someone blowing themselves up in a place that is crowded with civilians. Further supporting this goal is the observation that the coefficient for negative psychological or social impact is positive and reliable, indicating that this is a predetermination by the plotters of the suicide attack. Although damage to property may not be a prime objective, the greater the extent of this damage, the more we would expect to observe that it is from a suicide attack. We also note a complex relationship between the type of weapon that is used with the type of assailant, the type of attack and the rate of success. These data may not support clear distinctions of strategy with regard to weapons, except for perhaps bombs which are an integral part of almost all suicide attacks. In summary, the existence of a set of statistically reliable coefficient estimates in the presence of substantial heterogeneity in the actors studied shows that the GLMDM model can uncover common patterns by modelling the unexplained variance as a latent grouping.

The important question in an empirical study of terrorism *is not* whether we have explained the complete set of motivations for how and why these deadly attacks are developed; *it is* whether we have added anything to our current knowledge. This is because terrorism is such an important and vexing political problem. In this statistical application we coaxed some revealing information out of the data that increases understanding on the correlates of suicide attacks in the Middle East and northern Africa. Some of these findings are confirmatory, such as target preference and intergroup co-operation, whereas others, like the success and levels of damage that are inflicted by suicide attacks relative to non-suicide attacks, is surprising.

The results described above contain prescriptive advice for governmental organizations seeking to reduce the number and effectiveness of intentionally violent events. Although successful attacks are less associated with suicide strategies, suicide attackers generally inflict greater damage to property, additional injuries and, more importantly, more psychological or social impact

on the populace. Therefore countries in this region are justified in their counterterrorism policies that reduce training, remove motivation or interdict suicide attackers, even at the expense of less attention to non-suicide attackers. Strategic terrorism is mainly about attacking the social contract between a people and their government by eroding trust. The factors that are observed to have a stronger link to suicide attacks are more damaging to this relationship and therefore deserve more attention from governments that are concerned about the morale and support of their citizens.

Acknowledgements

Minjung Kyung, Jeff Gill and George Casella were supported by National Science Foundation grants DMS-0631632 and SES-0631588.

Appendix A: Fitting the logistic–Dirichlet model

A.1. Logistic distribution as a mixture of normals

The logistic distribution can be represented as a mixture of normals (see Andrews and Mallows (1974) or West (1987)), which leads us to a very efficient algorithm that can be put into a Gibbs sampler. If Z has a standard normal distribution, and $V/2$ has the asymptotic distribution of the Kolmogorov distance statistic and is independent of Z , then $Y = Z/V$ is logistic. From the identities in Andrews and Mallows (1974) (see also theorem 10.2.1 in Balakrishnan (1992)), for

$$f_X(x) = 8 \sum_{\alpha=1}^{\infty} (-1)^{\alpha+1} \alpha^2 x \exp(-2\alpha^2 x^2), \quad x \geq 0, \tag{9}$$

the KS density function (see Devroye (1986)), we can write

$$f_Y(y) = \int_0^{\infty} \left[\frac{1}{\sqrt{(2\pi)}} \frac{1}{2x} \exp\left\{-\frac{1}{2}\left(\frac{y}{2x}\right)^2\right\}\right] f_X(x) dx = \frac{\exp(-y)}{\{1 + \exp(-y)\}^2}, \tag{10}$$

where the density in square brackets is $N(0, 1/4x^2)$ and the result is the density function of a logistic distribution with mean 0 and variance $\pi^2/3$. Therefore, $Y \sim \mathcal{L}(0, \pi^2/3)$, where $\mathcal{L}(\cdot)$ denotes the logistic distribution.

Starting from the logistic distribution, we can model binary responses with a logit link function through a latent variable W_i such that

$$W_i = X_i\beta + \psi_i + \varepsilon_i, \quad \varepsilon_i \sim \mathcal{L}\left(0, \frac{\pi^2}{3}\sigma^2\right), \tag{11}$$

with $y_i = 1$ if $W_i > 0$ and $y_i = 0$ if $W_i \leq 0$, for $i = 1, \dots, n$. It can be shown that Y_i are independent Bernoulli random variables with $p_i = [1 + \exp\{-\mathbf{X}_i\beta - (\mathbf{A}\boldsymbol{\eta})_i\}]^{-1}$, the probability of success, and without loss of generality we fix $\sigma = 1$. With the mixture representation (10) an equivalent representation uses a normal latent variable, resulting in the likelihood function

$$\begin{aligned} L_k(\beta, \tau^2, \boldsymbol{\eta}, \mathbf{U}|\mathbf{A}, \mathbf{y}, \sigma^2) &= \prod_{i=1}^n \{I(U_i > 0)I(y_i = 1) + I(U_i \leq 0)I(y_i = 0)\} \\ &\times \int_0^{\infty} \left\{ \frac{1}{2\pi\sigma^2(2\xi)^2} \right\}^{n/2} \exp\left\{-\frac{1}{2\sigma^2(2\xi)^2}|\mathbf{U} - \mathbf{X}_i\beta - \mathbf{A}\boldsymbol{\eta}|^2\right\} \\ &\times 8 \sum_{\alpha=1}^{\infty} (-1)^{\alpha+1} \alpha^2 \xi \exp(-2\alpha^2\xi^2) d\xi \left(\frac{1}{2\pi\tau^2}\right)^{k/2} \exp\left(-\frac{1}{2\tau^2}|\boldsymbol{\eta}|^2\right), \end{aligned} \tag{12}$$

where $\mathbf{U} = (U_1, \dots, U_n)$, and U_i is the truncated normal variable such that

$$U_i = X_i\beta + (\mathbf{A}\boldsymbol{\eta})_i + \varepsilon_i, \quad \varepsilon_i \sim N\{0, \sigma^2(2\xi)^2\},$$

and $y_i = 1$ if $U_i > 0$ and $y_i = 0$ if $U_i \leq 0$, for $i = 1, \dots, n$.

A.2. Sampling schemes

An overview of the general sampling scheme is as follows. We identify three groups of parameters:

- (a) m , the precision parameter of the Dirichlet process,
- (b) \mathbf{A} , the indicator matrix of the partition defining the subclusters, and
- (c) $(\boldsymbol{\eta}, \boldsymbol{\beta}, \tau^2)$, the model parameters.

In the Gibbs sampler we iterate between these three groups until convergence.

Step 1: conditional on m and A , generate $(\boldsymbol{\eta}, \boldsymbol{\beta}, \tau^2) | \mathbf{A}, m$.

Step 2: conditional on $(\boldsymbol{\eta}, \boldsymbol{\beta}, \tau^2)$ and m , generate \mathbf{A} , a new partition matrix.

Step 3: conditional on $(\boldsymbol{\eta}, \boldsymbol{\beta}, \tau^2)$ and \mathbf{A} , generate m , the new precision parameter.

The model parameters are given the following prior distributions:

$$\left. \begin{aligned} \beta | \mu, \sigma^2 &\sim N(\mathbf{B}, d^* \sigma^2 I), \\ \mu &\sim \pi(\mu) \propto c \text{ (a flat prior with constant } c) \\ \tau^2 &\sim \text{inverted gamma}(a, b), \end{aligned} \right\} \tag{13}$$

where $\mathbf{B} = (\mu, 0, 0)'$, $d^* > 1$ and (a, b) are fixed such that the inverse gamma prior is diffuse ($a = 1$; b very small). We can either fix σ^2 or put a prior on it and estimate it in the hierarchical model with priors; here we shall fix a value for σ^2 . For the base measure of the Dirichlet process, we assume a normal distribution with mean 0 and variance τ^2 , $N(0, \tau^2)$.

A.2.1. Generating the logistic parameters

Given ξ , for fixed m and \mathbf{A} , a Gibbs sampler of $(\mu, \boldsymbol{\beta}, \tau^2, \boldsymbol{\eta}, \mathbf{U})$ is

$$\begin{aligned} \boldsymbol{\eta} | \mu, \boldsymbol{\beta}, \tau^2, \mathbf{U}, \mathbf{A}, \mathbf{y}, \sigma^2 &\sim N_k \left[\frac{1}{\sigma^2(2\xi)^2} \left\{ \frac{1}{\tau^2} I + \frac{1}{\sigma^2(2\xi)^2} \mathbf{A}'\mathbf{A} \right\}^{-1} \mathbf{A}'(\mathbf{U} - \mathbf{X}\boldsymbol{\beta}), \left\{ \frac{1}{\tau^2} I + \frac{1}{\sigma^2(2\xi)^2} \mathbf{A}'\mathbf{A} \right\}^{-1} \right], \\ \mu | \boldsymbol{\beta}, \tau^2, \boldsymbol{\eta}, \mathbf{U}, \mathbf{A}, \mathbf{y}, \sigma^2 &\sim N(\beta_0, d^* \sigma^2), \\ \boldsymbol{\beta} | \mu, \tau^2, \boldsymbol{\eta}, \mathbf{U}, \mathbf{A}, \mathbf{y}, \sigma^2 &\sim N_p \left[\left\{ \frac{1}{d^*} I + \frac{1}{(2\xi)^2} \mathbf{X}'\mathbf{X} \right\}^{-1} \left\{ \frac{1}{d^*} \mathbf{B} + \frac{1}{(2\xi)^2} \mathbf{X}'(\mathbf{U} - \mathbf{A}\boldsymbol{\eta}) \right\}, \right. \\ &\quad \left. \sigma^2 \left\{ \frac{1}{d^*} I + \frac{1}{(2\xi)^2} \mathbf{X}'\mathbf{X} \right\}^{-1} \right], \\ \tau^2 | \mu, \boldsymbol{\beta}, \boldsymbol{\eta}, \mathbf{U}, \mathbf{A}, \mathbf{y}, \sigma^2 &\sim \text{inverted gamma} \left(\frac{k}{2} + a, \frac{1}{2} |\boldsymbol{\eta}|^2 + b \right), \\ U_i | \boldsymbol{\beta}, \tau^2, \boldsymbol{\eta}, \mathbf{A}, y_i, \sigma^2 &\sim \begin{cases} N\{\mathbf{X}_i\boldsymbol{\beta} + (\mathbf{A}\boldsymbol{\eta})_i, \sigma^2(2\xi)^2\} & \text{if } y_i = 1, \\ N\{\mathbf{X}_i\boldsymbol{\beta} + (\mathbf{A}\boldsymbol{\eta})_i, \sigma^2(2\xi)^2\} & \text{if } y_i = 0 \end{cases} \end{aligned}$$

where $\mathbf{B} = (\mu, 0, 0)'$ and μ is from the posterior density. Then we update ξ from

$$\xi | \boldsymbol{\beta}, \tau^2, \boldsymbol{\eta}, \mathbf{U}, \mathbf{A}, \mathbf{y} \sim \left\{ \frac{1}{(2\xi)^2} \right\}^{n/2} \exp \left\{ -\frac{1}{2\sigma^2(2\xi)^2} |\mathbf{U} - \mathbf{X}\boldsymbol{\beta} - \mathbf{A}\boldsymbol{\eta}|^2 \right\} \times 8 \sum_{\alpha=1}^{\infty} (-1)^{\alpha+1} \alpha^2 \xi \exp(-2\alpha^2 \xi^2).$$

The conditional posterior density of ξ is the product of an inverted gamma prior with parameters $\alpha/2 - 1$ and $-(1/8\sigma^2)|\mathbf{U} - \mathbf{X}\boldsymbol{\beta} - \mathbf{A}\boldsymbol{\eta}|^2$, and the infinite sum of the sequence $(-1)^{\alpha+1} \alpha^2 \xi \exp(-2\alpha^2 \xi^2)$. To generate samples from this target density, we consider the alternating series method that was proposed by Devroye (1986). On the basis of his notation, we take

$$\begin{aligned} c h(\xi) &= 8 \left(\frac{1}{\xi^2} \right)^{n/2} \exp \left(-\frac{1}{8\sigma^2 \xi^2} |\mathbf{U} - \mathbf{X}\boldsymbol{\beta} - \mathbf{A}\boldsymbol{\eta}|^2 \right) \xi \exp(-2\xi^2), \\ a_n(\xi) &= (\alpha + 1)^2 \exp[-2\xi^2 \{(\alpha + 1)^2 - 1\}]. \end{aligned}$$

Here, we need to generate samples from $h(\xi)$, and we use accept-reject sampling with candidate $g(\xi^*) = 2 \exp(-2\xi^*)$, the exponential distribution with $\lambda = 2$, where $\xi^* = \xi^2$. Then we follow Devroye's method.

A.2.2. Dirichlet process parameters

In generating the Dirichlet process parameters we first update the partition matrix \mathbf{A} and then the precision parameter m . We use a Metropolis–Hastings algorithm with a candidate taken from a multinomial or Dirichlet distribution. This produces a Gibbs sampler that converges faster than the popular ‘stick breaking’ algorithm. (See Kyung *et al.* (2010b) for details.)

Given the model parameters at iteration t , $\theta^{(t)} = (\eta^{(t)}, \beta^{(t)}, \tau^{2(t)})$, we generate

$$\mathbf{q}^{(t+1)} = (q_1^{(t+1)}, \dots, q_n^{(t+1)}) \sim \text{Dirichlet}(n_1^{(t)} + 1, \dots, n_k^{(t)} + 1, 1, \dots, 1),$$

$$\mathbf{A}^{(t+1)} \sim P(\mathbf{A}') f(\mathbf{y}|\theta^{(t+1)}, \mathbf{A}') \binom{n}{n_1 \dots n_{k'}} \prod_{j=1}^{k'} (q_j^{(t+1)})^{n'_j} \tag{14}$$

where $n_j > 0, n_1 + \dots + n_k = n$.

Generating $\mathbf{q}^{(t+1)}$ is easy. To generate $\mathbf{A}^{(t+1)}$ we first generate a candidate that is an $n \times n$ matrix where each row is an independent multinomial distribution with probabilities $\mathbf{q}^{(t+1)}$, and the effective dimension of the matrix, the size of the partition, $k^{(t+1)}$, is the non-zero column sums. Deleting the columns with column sum 0 is a marginalization of the multinomial distribution. The probability of the candidate is given by

$$P(\mathbf{A}^{(t+1)}) = \frac{\Gamma(n)}{\Gamma(n - k^{(t+1)} + 1)} \frac{\Gamma(n_{k^{(t+1)}}^{(t+1)} + n - k^{(t+1)} + 1)}{\Gamma(2n)} \frac{k^{(t+1)-1}}{\prod_{j=1}^{k^{(t+1)}} \Gamma(n_j^{(t+1)} + 1)}$$

and a Metropolis–Hastings step is then done.

A.2.3. Generating the precision parameter

Rather than estimate m conventionally, a better strategy is to include m in the Gibbs sampler, as the marginal maximum likelihood estimate can be very unstable (Kyung *et al.*, 2010b). Conditional on $(\theta, \mathbf{A}, \mathbf{y})$, and k , with a prior $g(m)$ we obtain the posterior density

$$\pi(m|\theta, \mathbf{A}, \mathbf{y}) = \frac{\Gamma(m)}{\Gamma(m+n)} g(m)m^k \bigg/ \int_0^\infty \frac{\Gamma(m)}{\Gamma(m+n)} g(m)m^k dm, \tag{15}$$

where $\int \pi(m|\theta, \mathbf{A}, \mathbf{y}) dm < \infty$ as long as the exponent of m is positive. Note how far removed m is from the data, as the posterior depends only on the number of groups k . We consider a gamma distribution as a prior,

$$g(m) = \frac{m^{a-1} \exp(-m/b)}{\Gamma(a)b^a},$$

and generate m by using a Metropolis–Hastings algorithm with another gamma density as a candidate.

References

Abramowitz, M. and Stegun, I. A. (1972) Stirling numbers of the second kind. In *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, 9th printing, sect. 24.1.4, pp. 824–825. New York: Dover Publications.

Andrews, D. F. and Mallows, C. L. (1974) Scale mixtures of normal distributions. *J. R. Statist. Soc. B*, **36**, 99–102.

Antoniak, C. E. (1974) Mixtures of Dirichlet processes with applications to Bayesian nonparametric problems. *Ann. Statist.*, **2**, 1152–1174.

Arce, D. G. and Sandler, T. (2007) Terrorist signaling and the value of intelligence. *Br. J. Polit. Sci.*, **37**, 573–586.

Balakrishnan, N. (1992) *Handbook of the Logistic Distribution*. New York: CRC Press.

Blackwell, D. and MacQueen, J. B. (1973) Discreteness of Ferguson selections. *Ann. Statist.*, **1**, 355–358.

Breslow, N. E. and Clayton, D. G. (1993) Approximate inference in generalized linear mixed models. *J. Am. Statist. Ass.*, **88**, 9–25.

Bueno de Mesquita, E. (2005) Conciliation, counterterrorism, and patterns of terrorist violence. *Int. Organizn.*, **59**, 145–176.

Bueno de Mesquita, E. and Dickson, E. S. (2007) The propaganda of the deed: terrorism, counterterrorism, and mobilization. *Am. J. Polit. Sci.*, **51**, 364–381.

- Buonaccorsi, J. P. (1996) Measurement error in the response in the general linear model. *J. Am. Statist. Ass.*, **91**, 633–642.
- Carley, K. M. (2003) Dynamic network analysis. In *Dynamic Social Network Modeling and Analysis* (eds R. Breiger, K. Carley and P. Pattison). Washington DC: National Academies Press.
- Carley, K. M. (2004) Estimating vulnerabilities in large covert networks using multi-level data. In *Proc. North American Association for Computational Social and Organization Sciences Conf., Pittsburgh*. Pittsburgh: North American Association for Computational and Organization Sciences.
- Carley, K. (2006) A dynamic network approach to the assessment of terrorist groups and the impact of alternative courses of action. In *Proc. Visualizing Network Information*. Copenhagen: Royal Danish Defence College.
- Chai, S.-K. (1993) An organizational economics theory of antigovernment violence. *Compar. Polit.*, **26**, 99–110.
- Crenshaw, M. (1981) The causes of terrorism. *Compar. Polit.*, **13**, 379–399.
- Damien, P., Wakefield, J. and Walker, S. (1999) Gibbs sampling for Bayesian non-conjugate and hierarchical models by using auxiliary variables. *J. R. Statist. Soc. B*, **61**, 331–344.
- Devroye, L. (1986) *Non-uniform Random Variate Generation*. New York: Springer.
- Dey, D. K., Ghosh, S. K. and Mallick, B. K. (2000) *Generalized Linear Models: a Bayesian Perspective*. New York: Dekker.
- Dorazio, R. M., Mukherjee, B., Zhang, L., Ghosh, M., Jelks, H. L. and Jordan, F. (2007) Modelling unobserved sources of heterogeneity in animal abundance using a Dirichlet process prior. *Biometrics*, **64**, 635–644.
- Enders, W. (2007) Terrorism: an empirical analysis. In *Handbook of Defense Economics*, vol. 2 (eds K. Hartley and T. Sandler). Amsterdam: Elsevier.
- Enders, W. and Sandler, T. (1995) Terrorism: theory and applications. In *Handbook of Defense Economics*, vol. 1 (eds K. Hartley and T. Sandler). Amsterdam: Elsevier.
- Escobar, M. D. and West, M. (1995) Bayesian density estimation and inference using mixtures. *J. Am. Statist. Ass.*, **90**, 577–588.
- Fahrmeir, L. and Tutz, G. (2001) *Multivariate Statistical Modelling based on Generalized Linear Models*, 2nd edn. New York: Springer.
- Ferguson, T. S. (1973) A Bayesian analysis of some nonparametric problems. *Ann. Statist.*, **1**, 209–230.
- Gill, J. (2007) *Bayesian Methods for the Social and Behavioral Sciences*. New York: Chapman and Hall.
- Gill, J. and Casella, G. (2009) Nonparametric priors for ordinal Bayesian social science models: specification and estimation. *J. Am. Statist. Ass.*, **104**, 453–464.
- Jiang, J. (2007) *Linear and Generalized Linear Mixed Models and Their Applications*. New York: Springer.
- Krebs, V. E. (2002) Mapping terrorist cells. *Connections*, **24**, 43–52.
- Kydd, A. and Walter, B. F. (2002) Sabotaging the peace: the politics of extreme violence. *Int. Organizn*, **56**, 263–296.
- Kyung, M., Gill, J. and Casella, G. (2009) Characterizing the variance improvement in linear Dirichlet random effects models. *Statist. Probab. Lett.*, **79**, 2343–2350.
- Kyung, M., Gill, J. and Casella, G. (2010a) Estimation in Dirichlet random effects models. *Ann. Statist.*, **38**, 979–1009.
- Kyung, M., Gill, J. and Casella, G. (2010b) Sampling schemes for generalized linear Dirichlet process random effects models. *Statist. Meth. Applic.*, to be published.
- LaFree, G. and Dugan, L. (2008) *Global Terrorism Database II, 1998-2004*, computer file ICPSR22600-v2. Ann Arbor: Inter-university Consortium for Political and Social Research.
- Levy, S., Graham, H. D. and Gurr, T. R. (1969) A 150-year study of political violence in the United States. In *Violence in America: Historical and Comparative Perspectives (Official Report)*, pp. 81–91. Washington DC: National Commission on the Causes and Prevention of Violence.
- Li, Q. and Schaub, D. (2004) Economic globalization and transnational terrorist incidents. *J. Conflict Resoln*, **48**, 230–258.
- Little, R. J. A. and Rubin, D. B. (2002) *Statistical Analysis with Missing Data*, 2nd edn. New York: Wiley.
- Liu, J. S. (1996) Nonparametric hierarchical Bayes via sequential imputations. *Ann. Statist.*, **24**, 911–930.
- Lo, A. Y. (1984) On a class of Bayesian nonparametric estimates: I. density estimates. *Ann. Statist.*, **12**, 351–357.
- MacEachern, S. N. and Müller, P. (1998) Estimating mixture of Dirichlet process model. *J. Computnl Graph. Statist.*, **7**, 223–238.
- McCullagh, P. and Nelder, J. A. (1989) *Generalized Linear Models*, 2nd edn. New York: Chapman and Hall.
- McCullagh, P. and Yang, J. (2006) Stochastic classification models. In *Proc. Int. Congr. Mathematicians, Madrid, Aug. 22nd–30th*, vol. III, pp. 669–686. Helsinki: European Mathematical Society.
- McCulloch, C. E. and Searle, S. R. (2001) *Generalized, Linear, and Mixed Models*. New York: Wiley.
- Mickolus, E. F. (1982) *International Terrorism: Attributes of Terrorist Events, 1968-1977 (ITERATE 2)*. Ann Arbor: Inter-university Consortium for Political and Social Research.
- Mickolus, E. (2006) *International Terrorism: Attributes of Terrorist Events, 1968-2002*. Dunn Loring: Vinyard Software.
- Moon, I.-C. and Carley, K. M. (2007) Modeling and simulation of terrorist networks in social and geo-spatial dimensions. *IEEE Intell. Syst.*, **22**, 40–49.
- Neal, R. M. (2000) Markov chain sampling methods for Dirichlet process mixture models. *J. Computnl Graph. Statist.*, **9**, 249–265.

- Pape, R. A. (2006) *Dying to Win: Why Suicide Terrorists Do It*. London: Gibson Square Books.
- Rothenberg, R. (2002) From whole cloth: making up the terrorist network. *Connections*, **23**, 36–42.
- Sandler, T. and Arce, D. (2003) Terrorism and game theory. *Simuln Gamng*, **34**, 319–337.
- Siqueira, K. and Sandler, T. (2006) Terrorist versus the government: strategic interaction, support, and sponsorship. *J. Conflict Resoln*, **50**, 878–898.
- Teh, Y. W., Jordan, M. I., Beal, M. J. and Blei, D. M. (2006) Hierarchical Dirichlet processes. *J. Am. Statist. Ass.*, **101**, 1566–1581.
- Tsvetovat, M. and Carley, K. M. (2005) Structural knowledge and success of anti-terrorist activity: the downside of structural equivalence. *J. Socl Struct.*, **6**.
- Tsvetovat, M. and Carley, K. M. (2006) Improving effectiveness of communications sampling of covert networks. *E-Social Science Conf., Manchester*.
- Wang, N., Lin, X., Gutierrez, R. G. and Carroll, R. J. (1998) Bias analysis and SIMEX approach in generalized linear mixed measurement error models. *J. Am. Statist. Ass.*, **93**, 249–261.
- West, M. (1987) On scale mixtures of normal distributions. *Biometrika*, **74**, 646–648.
- Wolfinger, R. and O'Connell, M. (1993) Generalized linear mixed models: a pseudolikelihood approach. *J. Statist. Computn Simuln*, **48**, 233–243.